



Exotic carnivores in Flanders area expansion or repeated new input ?

Koen Van Den Berge & Jan Gouwy

koen.vandenberge@inbo.be & jan.gouwy@inbo.be



Since 1998, a volunteer network was installed by INBO, in co-operation with KBIN and funded by the Flemish Forest Administration, to collect victims of traffic among native Mustelids *Mustelidae* all over Flanders.

Place and date of dead specimen finds deliver basic elements for population distribution, but autopsy of the dead bodies generates additional information about their population ecology. In particular, the difference between erratic or settled animals is quite essential.

Collecting dead animals also allows a correct determination of the species – even when the body is very massacred or decomposed. E.g., the correct distinction between badgers *Meles meles* and other black-and-white rough-coated animals such as raccoon *Procyon lotor* and raccoon dog *Nyctereutes procyonoides* is not always evident, while American mink *Mustela vison* and polecat *Mustela putorius* can easily be confused.

These exotic carnivore species are believed to be on the point of colonizing Flanders for some decades. Thus, collecting dead mustelids was extended to all carnivores from the beginning of the research project.

Actually, raccoon, raccoon dog and American mink were indeed found as traffic casualties, beside some unexpected species such as striped skunk *Mephitis mephitis*, south American coati *Nasua nasua* and crab-eating raccoon *Procyon cancrivorus*.


For none of those animals however, autopsy revealed indications for reproduction in the wild. All of the specimen were most probably escaped animals in the first generation, mostly at very low numbers.

However, raccoon and – surprisingly – also striped skunk can be expected to establish wild populations from repeated escapes, as they seem both very popular pet animals.

Moreover, raccoon is actually establishing wild populations in the Walloon region at an accelerated tempo (personal observations), obviously as a 'logic result' of area expansion from the German border region. Reaching Flanders and mixing with escaped animals will only be a question of time.




Dead specimen collected from the wild since 1998



- *Procyon lotor* (13)
- *Procyon cancrivorus* (1)
- *Nyctereutes procyonoides* (2)
- *Mephitis mephitis* (4)
- *Mustela vison* (6)
- *Nasua nasua* (2)

For comparison : badger *Meles meles* 500
stone marten *Martes foina* 830
polecat *Mustela putorius* 1830



References

- Van Den Berge, K. (2008). Carnivore exoten in Vlaanderen. *Zoogdier* 19 (2) : 6-9
- Van Den Berge, K. & W. De Pauw (2003). Amerikaanse nerts *Mustela vison* (Schreber, 1777) / Wasbeer *Procyon lotor* (Linnaeus, 1758) / Wasbeerhond *Nyctereutes procyonoides* (Gray, 1834) . In: Verkem, S., De Maeseneer, J., Vandendriessche, B., Verbeyle, G. & Yskout, S. *Zoogdieren in Vlaanderen. Ecologie en verspreiding van 1987 tot 2002*. Natuurpunt Studie en JNM-Zoogdierenwerkgroep, Mechelen & Gent, België

Acknowledgements

We wish to thank Filip Berlengeer and Dirk Vansenant for their co-operation in the carnivore research and all the volunteers of the network for collecting dead carnivores